

CHAPTER 16: Pathogenic Bacteria Survival Through Cooking or Pasteurization

This guidance represents the Food and Drug Administration's (FDA's) current thinking on this topic. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. You can use an alternative approach if the approach satisfies the requirements of the applicable statutes and regulations. If you want to discuss an alternative approach, contact the FDA staff responsible for implementing this guidance. If you cannot identify the appropriate FDA staff, call the telephone number listed on the title page of this guidance.

UNDERSTAND THE POTENTIAL HAZARD.

The survival of pathogenic bacteria through cooking or pasteurization can cause consumer illness. The primary pathogens of concern are *Clostridium botulinum* (*C. botulinum*), *Listeria monocytogenes* (*L. monocytogenes*), *Campylobacter jejuni* (*C. jejuni*), pathogenic strains of *Escherichia coli* (*E. coli*), *Salmonella spp.*, *Shigella spp.*, *Yersinia enterocolitica* (*Y. enterocolitica*), *Staphylococcus aureus* (*S. aureus*), *Vibrio cholera* (*V. cholera*), *Vibrio vulnificus* (*V. vulnificus*), and *Vibrio parahaemolyticus* (*V. parahaemolyticus*). See Appendix 7 for a description of the public health impacts of these pathogens.

It is not practical to target viral pathogens in cooking or pasteurization processes because of their extreme heat resistance. Viral pathogens should be controlled through a rigorous sanitation regime as part of a prerequisite program or as part of Hazard Analysis Critical Control Point (HACCP) itself. The Procedures for the Safe and Sanitary Processing and Importing of Fish and Fishery Products regulation, 21 CFR 123 (called the Seafood HACCP Regulation in this guidance document) requires such a regime.

- **Types of heat processing**

Cooking is a heat treatment, usually performed before the product is placed in the finished product container. It is applied to fishery products that are distributed either refrigerated or frozen. Generally, after cooking, fishery products are referred to as cooked, ready to eat. Examples of cooked, ready-to-eat fishery products are crabmeat, lobster meat, crayfish meat, cooked shrimp, surimi-based analog products, seafood salads, seafood soups and sauces, and hot-smoked fish.

Pasteurization is a treatment (usually, but not always, the application of heat) applied to eliminate the most resistant pathogenic bacteria of public health concern that is reasonably likely to be present in the food for as long as the shelf-life of the product, when stored under normal and moderate abuse conditions. With fishery products, pasteurization is usually performed after the product is placed in the hermetically sealed finished product container. It is applied to fishery products that are distributed either refrigerated or frozen. Examples of pasteurized fishery products are pasteurized crabmeat, pasteurized surimi-based analog products, and pasteurized lobster meat.

In addition to eliminating bacterial pathogens, cooking and pasteurization also greatly reduce the number of spoilage bacteria present in the fishery product. These bacteria normally restrict the growth of pathogens through competition. Elimination of spoilage bacteria allows rapid growth of newly introduced pathogenic bacteria. Pathogenic bacteria that may be introduced after cooking or pasteurization are, therefore, a concern. This is especially true for pasteurization, because that process can significantly extend the shelf-life of the fishery product, providing more time for pathogenic bacteria growth and toxin formation.

Retorting is a heat treatment that eliminates all food-borne pathogens and produces a product that is shelf stable. Mandatory controls for retorting are provided in the Thermally Processed Low-Acid Foods Packaged in Hermetically Sealed Containers regulation, 21 CFR 113 (hereinafter, the Low Acid Canned Foods (LACF) Regulation), but are not covered in this chapter.

- **Goal of pasteurization**

Selection of the target pathogen is critical to the effectiveness of pasteurization. You should consider the potential that *C. botulinum* type E or non-proteolytic types B and F will survive the pasteurization process and grow under normal storage conditions or moderate abuse conditions. This is of particular concern if the product is reduced oxygen packaged (e.g., vacuum packaged or modified atmosphere packaged), does not contain a barrier that is sufficient to prevent growth and toxin formation by this pathogen, is not equipped with a time and temperature integrator, and is stored or distributed refrigerated (not frozen). In such products, you should ordinarily select *C. botulinum* type E and non-proteolytic types B and F as the target pathogen. For example, vacuum-packaged lobster meat that is pasteurized to kill *L. monocytogenes*, but not *C. botulinum* type E or non-proteolytic types B and F, and is not equipped with a Time-Temperature Indicator should be frozen to prevent growth and toxin formation by *C. botulinum* type E and non-proteolytic types B and F, and should be labeled to be held frozen and to be thawed under refrigeration immediately before use (e.g., “Important, keep frozen until used, thaw under refrigeration immediately before use”).

If the product is not reduced oxygen packaged, or contains a barrier that is sufficient to prevent the growth and toxin formation by *C. botulinum* type E or non-proteolytic types B and F, or is equipped with a time and temperature integrator, or is distributed frozen, then selection of another target pathogen may be appropriate. *L. monocytogenes* may be selected as the target pathogen for pasteurization of this type of product because it is the most resistant bacterial pathogen of public health concern that is reasonably likely to be present.

Surveys of retail display cases and home refrigerators indicate that temperatures above the minimum growth temperature of *C. botulinum* type E and non-proteolytic types B and F

(38°F (3.3°C)) are not uncommon. Therefore, refrigeration alone cannot be relied upon for control of the *C. botulinum* hazard. When freezing is relied upon to control the growth of *C. botulinum* type E and non-proteolytic types B and F, controls should be in place to ensure that the product is labeled with instructions that it be kept frozen throughout distribution.

For pasteurization processes that target *C. botulinum* type E and non-proteolytic types B and F, generally a reduction of six orders of magnitude (six logarithms, e.g., from 10^3 to 10^{-3}) in the level of contamination is suitable. This is called a 6D process. However, lower degrees of destruction may be acceptable if supported by a scientific study of the normal levels in the food before pasteurization. It is also possible that higher levels of destruction may be necessary in some foods, if especially high initial levels of the target pathogen are anticipated. Table A-4 (Appendix 4) provides 6D process times for a range of pasteurization temperatures, with *C. botulinum* type B (the most heat resistant form of non-proteolytic *C. botulinum*) as the target pathogen. The lethal rates and process times provided in the table may not be sufficient for the destruction of *C. botulinum* type E and non-proteolytic types B and F in dungeness crabmeat, because of the potential that naturally occurring substances, such as lysozyme, may enable the pathogen to more easily recover after heat damage.

Examples of properly pasteurized products are fish and fishery products generally (e.g., surimi-based products, soups, or sauces) pasteurized to a minimum cumulative total lethality of $F_{194^{\circ}\text{F}} (F_{90^{\circ}\text{C}}) = 10$ minutes, where $z = 12.6^{\circ}\text{F} (7^{\circ}\text{C})$ for temperatures less than 194°F (90°C) and $z = 18^{\circ}\text{F} (10^{\circ}\text{C})$ for temperatures above 194°F (90°C); blue crabmeat pasteurized to a minimum cumulative total lethality of $F_{185^{\circ}\text{F}} (F_{85^{\circ}\text{C}}) = 31$ minutes, where $z = 16^{\circ}\text{F} (9^{\circ}\text{C})$; and dungeness crabmeat pasteurized to a minimum cumulative total lethality of $F_{194^{\circ}\text{F}} (F_{90^{\circ}\text{C}}) = 57$ minutes, where $z = 15.5^{\circ}\text{F} (8.6^{\circ}\text{C})$. Equivalent processes at different temperatures can be calculated using the z values provided.

EXAMPLES OF PROPERLY PASTEURIZED PRODUCTS		
PRODUCT	MINIMUM CUMULATIVE TOTAL LETHALITY	Z VALUE
Fish and fishery products generally (e.g., surimi-based products, soups, or sauces)	$F_{194^{\circ}\text{F}} (F_{90^{\circ}\text{C}}) = 10$ minutes	12.6°F (7°C), for temperatures less than 194°F (90°C) 18°F (10°C) for temperatures above 194°F (90°C)
Blue crabmeat	$F_{185^{\circ}\text{F}} (F_{85^{\circ}\text{C}}) = 31$ minutes	16°F (9°C)
Dungeness crabmeat	$F_{194^{\circ}\text{F}} (F_{90^{\circ}\text{C}}) = 57$ minutes	15.5°F (8.6°C)

In some pasteurized surimi-based products, salt, in combination with a milder heat pasteurization process in the finished product container, works to prevent growth and toxin formation by *C. botulinum* type E and non-proteolytic types B and F. An example of a properly pasteurized surimi-based product in which 2.4% water phase salt is present is one that has been pasteurized at an internal temperature of 185°F (85°C) for at least 15 minutes. This process may not be suitable for other types of products because of the unique formulation and processing involved in the manufacture of surimi-based products.

Reduced oxygen-packaged foods that are pasteurized to control *C. botulinum* type E and non-proteolytic types B and F, but not *C. botulinum* type A and proteolytic types B and F, and that do not contain barriers to its growth should be refrigerated or frozen to control *C. botulinum* type A and proteolytic types B and F. Control of refrigeration is critical to the safety of these products. Further information on *C. botulinum* and reduced oxygen packaging is contained in Chapter 13.

In cases where *L. monocytogenes* is selected as the target pathogen, a 6D process is also generally suitable. FDA and U.S. Department of Agriculture's *L. monocytogenes* risk assessment indicates that approximately 8% of raw seafood are contaminated with from 1 to 10³ colony

forming unit (CFU)/g and that approximately 91% are contaminated at less than 1 CFU/g. Less than 1% of raw seafood are contaminated at levels greater than 10³ CFU/g and none at levels greater than 10⁶ CFU/g. FDA's limit for *L. monocytogenes* in ready-to-eat products, nondetectable, corresponds to a level of less than 1 CFU/25g.

Table A-3 (Appendix 4) provides 6D process times for a range of pasteurization temperatures, with *L. monocytogenes* as the target pathogen. Lower degrees of destruction may be acceptable if supported by a scientific study of the normal levels in the food before pasteurization. It is also possible that higher degrees of destruction may be necessary in some foods if especially high initial levels are anticipated.

Products that are pasteurized in the finished product container are at risk for recontamination after pasteurization. Controls, such as container seal integrity and protection from contaminated cooling water, are critical to the safety of these products and are covered in Chapter 18.

- **Goal of cooking for most products**

One reason for cooking products that will not be reduced oxygen packaged is to eliminate vegetative cells of pathogenic bacteria (or reduce them to an acceptable level) that may have been introduced to the process by raw materials or by processing that occurs before the cooking step. Selection of the target pathogen is critical to the effectiveness of cooking. Generally, *L. monocytogenes* is selected as the target pathogen because it is regarded as the most heat-tolerant, foodborne bacterial pathogen that does not form spores. Cooking processes are not usually designed to eliminate spores of bacterial pathogens. Determining the degree of destruction of the target pathogen is also critical. Generally, a reduction of six orders of magnitude (six logarithms, e.g., from 10³ to 10⁻³) in the level of contamination is suitable. This is called a 6D process.

Table A-3 provides 6D process times for a range of cooking temperatures, with *L. monocytogenes* as the target pathogen. Lower degrees of destruction

may be acceptable if supported by a scientific study of the normal levels in the food before pasteurization. It is also possible that higher degrees of destruction may be necessary in some foods if especially high initial levels are anticipated.

- **Goal of cooking refrigerated, reduced oxygen-packaged products**

Cooking is sometimes performed on products immediately before placement in reduced oxygen packaging (e.g., vacuum packaging or modified atmosphere packaging). These products include cooked, hot-filled soups, chowders, or sauces that are filled directly from the cook kettle using sanitary, automated, continuous filling systems designed to minimize risk of recontamination. They are often marketed under refrigeration, which is important for the control of *C. botulinum* type A and proteolytic types B and F.

The cooking process for these products should be sufficient to eliminate the spores of *C. botulinum* type E and non-proteolytic types B and F. This is the case when the product does not contain other barriers that are sufficient to prevent growth and toxin formation by this pathogen. Generally, a 6D process (six logarithms, e.g., from 10^3 to 10^{-3}) is suitable. However, lower degrees of destruction may be acceptable if supported by a scientific study of the normal levels in the food before pasteurization. It is also possible that higher degrees of destruction may be necessary in some foods if especially high initial levels are anticipated.

Table A-4 provides 6D process times for a range of cooking temperatures, with *C. botulinum* type B (the most heat-resistant form of non-proteolytic *C. botulinum*) as the target pathogen. The lethal rates and process times provided in the table may not be sufficient for the destruction of *C. botulinum* type E and non-proteolytic types B and F in soups or sauces containing dungeness crabmeat because of the potential that naturally occurring substances, such as lysozyme, may enable the pathogen to more easily recover after damage. An example of a product that is

properly cooked to eliminate *C. botulinum* type E and non-proteolytic types B and F is a soup or sauce that is cooked to a minimum cumulative total lethality of $F_{194^{\circ}\text{F}} (F_{90^{\circ}\text{C}}) = 10$ minutes, where $z = 12.6^{\circ}\text{F} (7^{\circ}\text{C})$ for temperatures less than $194^{\circ}\text{F} (90^{\circ}\text{C})$ and $z = 18^{\circ}\text{F} (10^{\circ}\text{C})$ for temperatures above $194^{\circ}\text{F} (90^{\circ}\text{C})$.

Reduced oxygen-packaged soups or sauces that are cooked immediately before packaging to control *C. botulinum* type E and non-proteolytic types B and F, but not *C. botulinum* type A and proteolytic types B and F, and that do not contain barriers to its growth should be refrigerated or frozen to control *C. botulinum* type A and proteolytic types B and F. Control of refrigeration is critical to the safety of these products. Further information on *C. botulinum* and reduced oxygen packaging is contained in Chapter 13.

Cooking processes that target *C. botulinum* type E and non-proteolytic types B and F have much in common with pasteurization processes. Like products that are pasteurized in the final container, products that are cooked and then placed in the final container also are at risk for recontamination after they are placed in the finished product container. Controls, such as container seal integrity and protection from contaminated cooling water, are critical to the safety of these products and are covered in Chapter 18.

Additionally, because these products are cooked before they are packaged, they are at risk of recontamination between cooking and packaging. The risk of recontamination may be minimized by filling the container in a sanitary, automated, continuous filling system while the product is still hot (i.e., hot filling). This is another critical step for the safety of these products. This control strategy is suitable for products that are filled directly from the cooking kettle, where the risk of recontamination is minimized. It is not ordinarily suitable for products such as crabmeat, lobster meat, or crayfish meat that are handled between cooking and filling. Hot filling is also covered in Chapter 18.

- **Control by cooking or pasteurization**

Controlling pathogenic bacteria survival through cooking or pasteurization is accomplished by:

- Scientifically establishing a cooking or pasteurization process that will eliminate pathogenic bacteria of public health concern or reduce their numbers to acceptable levels;
- Designing and operating the cooking or pasteurization equipment so that every unit of product receives at least the established minimum process;
- Continuously monitoring the critical process parameters to verify achievement of a scientifically established process (e.g., time and temperature).

You may monitor End-Point Internal Product Temperature (EPIPT), a measurement of the temperature of the product as it exits the heat process, instead of performing continuous time and temperature monitoring. This approach is suitable if you have conducted a scientific study to validate that the EPIPT that you have selected will provide an appropriate reduction in the numbers of the target pathogen (e.g., 6D) in the slowest heating unit or portion of product under the worst set of heating conditions covered by the scientific study. You should (1) conduct a temperature distribution study within the heating system to identify any cold spots; (2) conduct a heat penetration study that accounts for the slowest heating product under the worst case heating conditions covered by the scientific study; and identify other critical factors of processing and/or packaging that affect the rate of product heating when scientifically establishing a cooking or pasteurization process (i.e., process validation). The EPIPT should be used as a monitoring technique only under those conditions that were evaluated by the scientific study. Those conditions may need to be identified as critical limits and monitored as part of the HACCP plan.

EPIPT monitoring may not be an option when the objective is control of *C. botulinum* type E and non-proteolytic types B and F spores. These spores are far more heat resistant than vegetative

cells of *L. monocytogenes* and destroying them requires an EPIPT that could be achieved only in a pressurized steam environment, making measurement impractical. Additional guidance on EPIPT monitoring can be found in Food Processors Association guidance document “FPA Guidance Document: Establishing or Verifying a Heat Process for Cooked, Ready-to-Eat Seafood Products, and Heat Process Monitoring Considerations under HACCP,” 2nd Edition, February 2005 and purchased at the Grocery Manufacturers Association, Washington DC 20005.

- **Strategies for controlling pathogenic bacteria growth**

There are a number of strategies for the control of pathogenic bacteria in fish and fishery products. They include:

- Killing pathogenic bacteria by cooking or pasteurizing (covered in this chapter) or retorting (covered by the LACF Regulation, 21 CFR 113);
- Killing pathogenic bacteria by processes that retain the raw characteristics of the products (covered in Chapter 17);
- Managing the amount of time that food is exposed to temperatures that are favorable for pathogenic bacteria growth and toxin production (covered generally in Chapter 12; for *C. botulinum*, in Chapter 13; and for *S. aureus* in hydrated batter mixes, in Chapter 15);
- Controlling the amount of moisture that is available for pathogenic bacteria growth (water activity) in the product by drying (covered in Chapter 14);
- Controlling the amount of moisture that is available for pathogenic bacteria growth (water activity) in the product by formulation (covered in Chapter 13);
- Controlling the amount of salt or preservatives, such as sodium nitrite, in the product (covered in Chapter 13);
- Controlling the level of acidity (pH) in the product (covered by the Acidified Foods

regulation, 21 CFR 114, for shelf-stable acidified products, and by Chapter 13 for refrigerated acidified products);

- Controlling the source of molluscan shellfish and the time from exposure to air (e.g., by harvest or receding tide) to refrigeration to control pathogens from the harvest area (covered in Chapter 4);
- Controlling the introduction of pathogenic bacteria after the pasteurization process (covered in Chapter 18).

DETERMINE WHETHER THE POTENTIAL HAZARD IS SIGNIFICANT.

The following guidance will assist you in determining whether pathogenic bacteria survival through cooking and pasteurization is a significant hazard at a processing step.

1. Is it reasonably likely that unsafe levels of pathogenic bacteria will be introduced at this processing step (do unsafe levels of pathogenic bacteria come in with the raw material, or will the process introduce unsafe levels of pathogenic bacteria)?

It is reasonable to assume that pathogens of various types, including those listed in Table A-1 (Appendix 4), will be present on raw fish and fishery products. They may be present only at low levels or only occasionally, but even such occurrences warrant consideration because of the potential for growth and toxin production.

Pathogenic bacteria may also be introduced during processing, from the air, unclean hands, insanitary utensils and equipment, unsafe water, and sewage. Well-designed sanitation programs will minimize the introduction of pathogens. Such sanitation controls need not be part of your HACCP plan if they are monitored under your sanitation program (prerequisite program). In most cases, it is not reasonable to assume that they will fully prevent the introduction

of bacterial pathogens. For this reason, you should consider it reasonably likely that low numbers of pathogenic bacteria will be present in the product.

2. Can unsafe levels of pathogenic bacteria that were introduced at an earlier processing step be eliminated or reduced to an acceptable level at this processing step?

Pathogenic bacteria survival through cooking or pasteurization should also be considered a significant hazard at any processing step where a preventive measure is, or can be, used to eliminate the hazard (or reduce the likelihood of its occurrence to an acceptable level) if it is reasonably likely to occur. The preventive measure that can be applied for pathogenic bacteria survival through cooking and pasteurization is proper design and control of the cooking or pasteurization process.

- **Intended use**

Because cooked or pasteurized products are ready to eat, it is unlikely that the intended use will affect the significance of the hazard.

IDENTIFY CRITICAL CONTROL POINTS.

The following guidance will assist you in determining whether a processing step is a critical control point (CCP) for the survival of pathogenic bacteria through cooking or pasteurization:

Will the finished product be pasteurized in the final container?

1. If the finished product will be pasteurized in the final container, you should identify the pasteurization step as the CCP. In this case, you would not need to identify the cooking step as a CCP for the hazard of pathogenic bacteria survival through cooking.

Example:

A crabmeat processor cooks, picks, packs, and pasteurizes the crabmeat.

The processor sets the CCP for pathogenic bacteria survival through cooking and pasteurization at the pasteurization step and does not identify the cooking step as a CCP for this hazard.

This control approach is a control strategy referred to in this chapter as “Control Strategy Example - Cooking and Pasteurization.”

2. If the product will not be pasteurized, you should identify the cooking step as the CCP.

This control approach is the same as the one above and is a control strategy also referred to in this chapter as “Control Strategy Example - Cooking and Pasteurization.” For products in reduced oxygen packaging for which the cooking process does not target *C. botulinum* type E and non-proteolytic types B and F, see Chapter 13 for additional guidance.

DEVELOP A CONTROL STRATEGY.

The following guidance provides a control strategy for survival of pathogenic bacteria through cooking or pasteurization. You may select a control strategy that is different from that which is suggested, provided it complies with the requirements of the applicable food safety laws and regulations.

The following is an example of the control strategy included in this chapter:

CONTROL STRATEGY	MAY APPLY TO PRIMARY PROCESSOR	MAY APPLY TO SECONDARY PROCESSOR
Cooking and pasteurization	✓	✓

• CONTROL STRATEGY EXAMPLE - COOKING AND PASTEURIZATION

Set Critical Limits.

- The minimum or maximum values for the critical factors established by a scientific study. These may include length of the

cook or pasteurization cycle (speed of the belt for a continuous cooker or pasteurizer), temperature of the steam or water used for cooking or pasteurization (or visual observation of minutes at a boil for cooking), initial temperature of the product, container size (e.g., can dimensions, pouch thickness), and product formulation. Other critical factors that affect the rate of heating of the product may also be established by the study;

OR

- The EPIPT, established by a scientific study. Other critical factors that affect the rate of heating of the product may also be established by the study.

Note: EPIPT monitoring may not be an option when the objective is control of *C. botulinum* type E and non-proteolytic types B and F spores.

Establish Monitoring Procedures.

» What Will Be Monitored?

- The critical factors established by a scientific study. These may include length of the cook or pasteurization cycle (speed of the belt for a continuous cooker or pasteurizer) and temperature of the steam or water used for cooking or pasteurization (or visual observation of minutes at a boil for cooking), initial temperature of the product, container size (e.g., can dimensions, pouch thickness), and product formulation;

OR

- The EPIPT.

» How Will Monitoring Be Done?

For batch cooking or pasteurization equipment:

- For cooking or pasteurization temperature:
 - Use a continuous temperature-recording device (e.g., a recording thermometer). The device should be installed where it measures the coldest temperature of the cooking equipment (cold spot to be determined by a study). Where cooking

is performed at the boiling point, visual observation of minutes at a boil may be an acceptable alternative;

AND

- For the start and end of each cooking or pasteurization cycle:
 - Visual observation;

AND

- For other critical factors:
 - Use equipment appropriate to the critical factor (e.g., initial temperature with a temperature-indicating device, (e.g., a thermometer);

OR

- For the EPIPT:
 - Use a temperature-indicating device (e.g., a thermometer).

For continuous cooking or pasteurization equipment:

- For cooking or pasteurization temperature:
 - Use a continuous temperature-recording device (e.g., a recording thermometer). The device should be installed where it measures the coldest temperature of the cooking equipment (cold spot to be determined by a study). Because of the extended time of operation of such equipment, it is unlikely that visual observation of boiling will be an acceptable alternative, even if cooking is performed at the boiling point;

AND

- For cooking or pasteurization time, use:
 - A stopwatch or tachometer to monitor the speed of the belt drive wheel;

OR

- A stopwatch to monitor the time necessary for a test unit or belt marking to pass through the equipment;

AND

- For other critical factors:

- Use equipment appropriate to the critical factor (e.g., initial temperature with a temperature-indicating device, (e.g., a thermometer);

OR

- For the EPIPT:
 - Use a temperature-indicating device (e.g., a thermometer).

» **How Often Will Monitoring Be Done (Frequency)?**

For batch cooking or pasteurization equipment:

- For cooking or pasteurization temperature:
 - Continuous monitoring, with a visual check of the recorded data at least once per batch;

AND

- For the start and end of each cooking or pasteurization cycle:
 - Each batch;

AND

- For other critical factors:
 - With sufficient frequency to achieve control;

OR

- For the EPIPT:
 - Each batch.

For continuous cooking or pasteurization equipment:

- For cooking or pasteurization temperature:
 - Continuous monitoring, with a visual check of the recorded data at least once per day;

AND

- For cooking or pasteurization time:
 - At least once per day, and whenever any changes in belt speed are made;

AND

- For other critical factors:
 - With sufficient frequency to achieve control;

OR

- For the EPIPT:
 - At least every 30 minutes, and whenever any changes in product-heating critical factors occur.

» **Who Will Perform the Monitoring?**

- For continuous temperature-recording devices:
 - Monitoring is performed by the device itself. The visual check of the data generated by the device, to ensure that the critical limits have consistently been met, may be performed by any person who has an understanding of the nature of the controls;

AND

- For other monitoring:
 - Any person who has an understanding of the nature of the controls.

Establish Corrective Action Procedures.

Take the following corrective action to a product involved in a critical limit deviation:

- Recook or repasteurize the product;
- OR
- Chill and hold the product for an evaluation of the adequacy of the cooking or pasteurization process. If the product has not received an adequate process, it should be destroyed, diverted to a non-food use, or recooked or repasteurized;
- OR
- Divert the product to a use in which the critical limit is not applicable (e.g., divert improperly cooked or pasteurized shrimp to a shrimp canning operation);
- OR
- Destroy the product;
- OR
- Divert the product to a non-food use.

AND

Take the following corrective action to regain control over the operation after a critical limit deviation:

- Adjust the steam supply to increase the processing temperature;
- OR
- Extend the length of the cooking or pasteurization cycle to compensate for a temperature drop, using a process developed by a process authority;
- OR
- Process at a higher temperature to compensate for a low initial temperature, using a process developed by a process authority;
- OR
- Adjust the belt speed.

Establish a Recordkeeping System.

For batch cooking or pasteurization equipment:

- For temperature monitoring:
 - Record of continuous temperature monitoring;
- AND
- Record of visual checks of recorded data;
- OR
- Cooking log that indicates visual observation of boiling, where cooking is performed at the boiling point;
- AND
- Record of notation of the start time and end time of the cooking or pasteurization periods;
- AND
- Records that are appropriate for the other critical factors (e.g., a cooking or pasteurization log that indicates the initial temperature);
- OR
- Record of EPIPT results.

For continuous cooking or pasteurization equipment:

- Record of continuous temperature monitoring;
AND
- Record of visual checks of devices;
AND
- Cooking or pasteurization log that indicates the RPM of the belt drive wheel or the time necessary for a test unit or belt marking to pass through the tank;
AND
- Records that are appropriate for the other critical factors (e.g., a cooking or pasteurization log that indicates the initial temperature);
OR
- Record of EPIPT results.

Establish Verification Procedures.

For cooking, process validation study (process establishment):

- The adequacy of the cooking process should be established by a scientific study. It should be designed to ensure an appropriate reduction in the number of pathogenic bacteria of public health concern. Selecting the target organism is critical. In most cases, it will be a relatively heat-tolerant vegetative pathogen, such as *L. monocytogenes*. However, in some cases where outgrowth of spore-forming pathogens, such as *Clostridium perfringens* and *Bacillus cereus*, during the post-cook cooling step must be prevented by eliminating these pathogens during the cook step (e.g., because cooling after cooking is not controlled (see Chapter 12)), then they will be the target organisms. Additionally, when cooking is performed immediately before reduced oxygen packaging (e.g., vacuum packaging or modified atmosphere packaging), for a product that will be marketed under refrigeration, it may be necessary for the cooking process to be sufficient to eliminate

the spores of *C. botulinum* type E and non-proteolytic types B and F. This is the case when the product does not contain other barriers that are sufficient to prevent growth and toxin formation by this pathogen (e.g., refrigerated, vacuum packaged hot-filled soups and sauces). Generally, a 6D process is suitable, regardless of the target bacterial pathogen. However, lower degrees of destruction may be acceptable if supported by a scientific study of the normal levels in the food. Tables A-3 and A-4 provide 6D process times for a range of internal product temperatures, with *L. monocytogenes* and *C. botulinum* type B (the most heat-resistant form of non-proteolytic *C. botulinum*) as the target pathogens. The values provided in Table A-4 may not be sufficient for the destruction of *C. botulinum* type E and non-proteolytic types B and F in products containing dungeness crabmeat because of the potential protective effect of naturally occurring substances, such as lysozyme.

Expert knowledge of thermal process calculations and the dynamics of heat transfer in processing equipment may be required to establish such a cooking process. Such knowledge can be obtained by education or experience, or both. Conducting a validation study for cooking processes may require access to suitable facilities and the application of recognized methods. The cooking equipment should be designed, operated, and maintained to deliver the established process to every unit of the product. In some cases, thermal death time, heat penetration, temperature distribution, and inoculated pack studies may be necessary to validate the minimum process. In many cases, establishing the minimum process may be simplified by repetitively determining the process needed to reach an internal product temperature that will ensure the inactivation of all vegetative bacterial pathogens of public health concern under the most difficult heating conditions likely to be encountered

during processing. In other instances, existing literature or federal, state, or local regulations that establish minimum processes or adequacy of equipment are available. Characteristics of the process, product, and/or equipment that affect the ability of the established minimum cooking process should be taken into consideration in the validation of the process. A record of the process validation study should be maintained;

OR

For pasteurization, process validation study (process establishment):

- The adequacy of the pasteurization process should be established by a scientific study. It should be designed to ensure an appropriate reduction in the number of target bacterial pathogens. Selecting the target organism is critical. In most cases, it will be the spores of *C. botulinum* type E and non-proteolytic types B and F. In some cases (e.g., products that are distributed frozen or contain other barriers to prevent growth and toxin formation by *C. botulinum* type E and non-proteolytic types B and F), the process will target another pathogen, such as *L. monocytogenes*. Generally, a 6D process is suitable, regardless of the target pathogen. However, lower degrees of destruction may be acceptable if supported by a scientific study of the normal levels in the food. Tables A-3 and A-4 provide 6D process times for a range of internal product temperatures, with *L. monocytogenes* and *C. botulinum* type B (the most heat-resistant form of non-proteolytic *C. botulinum*) as the target pathogens. The values provided in Table A-4 may not be sufficient for the destruction of *C. botulinum* type E and non-proteolytic types B and F in products containing dungeness crabmeat because of the potential protective effect of naturally occurring substances, such as lysozyme.

Expert knowledge of thermal process calculations and the dynamics of heat transfer

in processing equipment may be required to determine the target bacterial pathogen and to establish a pasteurization process. Such knowledge can be obtained by education or experience, or both. Conducting a validation study for pasteurization processes may require access to suitable facilities and the application of recognized methods. The pasteurization equipment should be designed, operated, and maintained to deliver the established process to every unit of the product. In some cases, thermal death time, heat penetration, temperature distribution, and inoculated pack studies may be necessary to validate the minimum process. In other instances, existing literature or federal, state, or local regulations that establish minimum processes or adequacy of equipment are available. Characteristics of the process, product, and/or equipment that affect the adequacy of the established minimum pasteurization process should be taken into consideration in the validation of the process. A record of the validation study should be maintained;

AND

- Before a temperature-indicating device (e.g., a thermometer) or temperature-recording device (e.g., a recording thermometer) is put into service, check the accuracy of the device to verify that the factory calibration has not been affected. This check can be accomplished by:
 - Immersing the sensor in an ice slurry (32°F (0°C)) if the device will be used at or near refrigeration temperature;
- OR
- Immersing the sensor in boiling water (212°F (100°C)) if the device will be used at or near the boiling point (note that the temperature should be adjusted to compensate for altitude, when necessary);
- OR
- A combination of the above if the

device will be used at or near room temperature;

OR

- Comparing the temperature reading on the device with the reading on a known accurate reference device (e.g., a thermometer traceable to National Institute of Standards and Technology (NIST) standards) under conditions that are similar to how it will be used (e.g., steam temperature, water temperature, product internal temperature) within the temperature range at which it will be used;

AND

- Once in service, check the temperature-indicating device or temperature-recording device daily before the beginning of operations. Less frequent accuracy checks may be appropriate if they are recommended by the instrument manufacturer and the history of use of the instrument in your facility has shown that the instrument consistently remains accurate for a longer period of time. In addition to checking that the device is accurate by one of the methods described above, this process should include a visual examination of the sensor and any attached wires for damage or kinks. The device should be checked to ensure that it is operational and, where applicable, has sufficient ink and paper;

AND

- Calibrate the temperature-indicating device or temperature-recording device against a known accurate reference device (e.g., a NIST-traceable thermometer) at least once a year or more frequently if recommended by the device manufacturer. Optimal calibration frequency is dependent upon the type, condition, past performance, and conditions of use of the device. Consistent temperature variations away from the actual value (drift) found during checks and/or calibration may

show a need for more frequent calibration or the need to replace the device (perhaps with a more durable device). Devices subjected to high temperatures for extended periods of time may require more frequent calibration. Calibration should be performed at a minimum of two temperatures that bracket the temperature range at which it is used;

AND

- Calibrate other instruments as necessary to ensure their accuracy;

AND

- Review monitoring, corrective action, and verification records within 1 week of preparation to ensure they are complete and any critical limit deviations that occurred were appropriately addressed.

TABLE 16-1

**CONTROL STRATEGY EXAMPLE - COOKING AND PASTEURIZATION
(COOKING MODEL)**

This table is an example of a portion of a HACCP plan using "Control Strategy Example - Cooking and Pasteurization (Cooking Model)." This example illustrates how a processor of wild-caught cooked shrimp can control cooking using a continuous steam cooker. It is provided for illustrative purposes only.

Pathogenic bacteria survival through cooking and pasteurization may be only one of several significant hazards for this product. Refer to Tables 3-3 and 3-4 (Chapter 3) for other potential hazards (e.g., environmental chemical contaminants and pesticides, pathogenic bacteria growth and toxin formation during processing, food and color additives, and metal fragments).

**Example Only
See Text for Full Recommendations**

(1)	(2)	(3)	(4)	(5)			(7)	(8)	(9)	(10)
				WHAT	HOW	FREQUENCY				
Cooking	Pathogenic bacteria survival	Minimum cook time: 2.5 minutes Minimum cook temperature: 210°F Note: To achieve a 6D reduction of L. monocytogenes	Length of the cook cycle	Belt speed measurement with stopwatch	Once per day and after any adjustment	Cooker operator	Extend process or elevate temperature to compensate for deviation from critical limit, based on alternate processes provided by the process authority Chill and hold for evaluation	Cooking record Data logger printout Grading record	Scientific study establishing the thermal process (process validation) Check the data logger for accuracy and damage and to ensure that it is operational before putting into operation; check it daily, at the beginning of operations; and calibrate it once per year Calibrate the scale monthly Review monitoring, corrective action and verification, records within 1 week of preparation	
				Digital time and temperature data logger	Continuous, with visual check of recorded data once per day	Cooker operator				
				Scale	Hourly and after every raw material lot change or grader adjustment	Grader operator				

*Note: The critical limits in this example are for illustrative purposes only and are not related to any recommended process.

TABLE 16-2

**CONTROL STRATEGY EXAMPLE - COOKING AND PASTEURIZATION
(PASTEURIZATION MODEL)**

This table is an example of a portion of a HACCP plan using “Control Strategy Example - Cooking and Pasteurization (Pasteurization Model).” This example illustrates how a processor of pasteurized, refrigerated blue crabmeat can control pasteurization. It is provided for illustrative purposes only.

Pathogenic bacteria survival through cooking and pasteurization may be only one of several significant hazards for this product. Refer to Tables 3-3 and 3-4 (Chapter 3) for other potential hazards (e.g., environmental chemical contaminants and pesticides, pathogenic bacteria growth and toxin formation during processing, recontamination after pasteurization, and metal fragments).

**Example Only
See Text for Full Recommendations**

(1)	(2)	(3)	(4)	(5)			(7)	(8)	(9)	(10)
				WHAT	HOW	FREQUENCY				
Batch pasteurization	Pathogenic bacteria survival	CRITICAL LIMITS FOR EACH PREVENTIVE MEASURE* Minimum initial product temperature: 37°F Minimum length of pasteurization cycle: 120 minutes	Initial temperature	Dial thermometer	Coldest can entering each batch	Pasteurizer operator	Extend the process or elevate the temperature to compensate for deviation from the critical limit Segregate and hold for evaluation	Pasteurization log	Process establishment Check the temperature-recording device and dial thermometer for accuracy and damage and to ensure that they are operational before putting into operation; check it daily, at the beginning of operations; and calibrate it once per year Review monitoring, verification, and corrective action records within 1 week of preparation	
			Time up to 189°F and time cycle ends	Temperature-recording device	Each batch	Pasteurizer operator		Pasteurization log		
			Temperature of water bath	Temperature-recording device	Continuously, with visual check at end of batch	Recorder thermometer, with visual check by pasteurizer operator		Recorder thermometer chart		

*Note: The critical limits in this example are for illustrative purposes only and are not related to any recommended process.

BIBLIOGRAPHY.

We have placed the following references on display in the Division of Dockets Management, Food and Drug Administration, 5630 Fishers Lane, rm. 1061, Rockville, MD 20852. You may see them at that location between 9 a.m. and 4 p.m., Monday through Friday. As of March 29, 2011, FDA had verified the Web site address for the references it makes available as hyperlinks from the Internet copy of this guidance, but FDA is not responsible for any subsequent changes to Non-FDA Web site references after March 29, 2011.

- Cockey, R. R., and M. C. Tatro. 1974. Survival studies with spores of *Clostridium botulinum* type E in pasteurized meat of the blue crab *Callinectes sapidus*. *Appl. Microbiol.* 27:629-633.
- European Chilled Food Federation. 1997. Guidelines for good hygienic practice in the manufacture of chilled foods.
- Frazier, J. 2005. Establishing or verifying a heat process for cooked, ready-to-eat seafood products, and heat process monitoring considerations under HACCP. 2nd ed. Grocery Manufacturers Association (Food Products Association), Washington, DC.
- Hilderbrand, K. S., Jr. 1996. Personal communication. Oregon State University, Extension Service, Corvallis, OR.
- Lum, K. C. 1996. Personal communication. National Food Processors Association, Seattle, WA.
- Lynt, R. K., D. A. Kautter, and H. M. Solomon. 1982. Differences and similarities among proteolytic and nonproteolytic strains of *Clostridium botulinum* types A, B, E and F: a review. *J. Food Prot.* 45:466-474.
- Lynt, R. K., H. M. Solomon, T. Lilly, and D. A. Kautter. 1977. Thermal death time of *Clostridium botulinum* type E in meat of the blue crab. *J. Food Sci.* 42:1022-1025.
- Mackey, B. M., and N. Bratchell. 1989. The heat resistance of *Listeria monocytogenes*: a review. *Lett. Appl. Microbiol.* 9:89-94.
- National Advisory Committee on Microbiological Criteria for Foods. 1990. Recommendations of the National Advisory Committee on Microbiological Criteria for Foods for Cooked Ready-to-Eat Shrimp and Cooked Ready-to-Eat Crabmeat. Executive Secretariat, Food Safety and Inspection Service, U.S. Department of Agriculture, Washington, DC.
- National Advisory Committee on Microbiological Criteria for Foods. 1990. Recommendations of the National Advisory Committee on Microbiological Criteria for Foods for Refrigerated Foods Containing Cooked, Uncured Meat or Poultry Products that are Packaged for Extended Refrigerated Shelf Life and that are Ready-to-Eat or Prepared with Little or No Additional Heat Treatment. Executive Secretariat, Food Safety and Inspection Service, U.S. Department of Agriculture, Washington, DC.
- National Advisory Committee on Microbiological Criteria for Foods. 1991. *Listeria monocytogenes*: recommendations of the National Advisory Committee on Microbiological Criteria for Foods for Refrigerated Foods. *Intl. J. Food Microbiol.* 14:185-246.
- Peterson, M. E., G. A. Pelroy, F. T. Poysky, R. N. Paranjpye, R. M. Dong, G. M. Pigott, and M. W. Eklund. 1997. Heat-pasteurization process for inactivation of nonproteolytic types of *Clostridium botulinum* in picked dungeness crabmeat. *J. Food Prot.* 60:928-934.
- Peterson, M. E., R. N. Paranjpye, F. T. Poysky, G. A. Pelroy, and M. W. Eklund. 2002. Control of nonproteolytic *Clostridium botulinum* types B and E in crab analogs by combinations of heat pasteurization and water phase salt. *J. Food Prot.* 65:130-139.
- Rippen, T., C. Hackney, G. Flick, G. Knobl, and D. Ward. 1993. Seafood pasteurization and minimal processing manual. Virginia Cooperative Extension Publication 600-0061. Virginia Sea Grant Publication VSG

93-09. Virginia Seafood Research and Extension Center Publication VPI-SG-93-01. Blacksburg, VA.

- U.S. Food and Drug Administration and U.S. Department of Agriculture. 2003. Quantitative assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. <http://www.fda.gov/Food/ScienceResearch/ResearchAreas/RiskAssessmentSafetyAssessment/ucm183966.htm>.
- U.S. Food and Drug Administration. Thermally processed low-acid foods packaged in hermetically sealed containers. *In* Code of Federal Regulations, 21 CFR 113. U.S. Government Printing Office, Washington, DC.